

Late Preterm Infants

Morbidities, Mortality, and Management Recommendations

Katie Huff, MD, Rebecca S. Rose, MD, William A. Engle, MD*

KEYWORDS

• Late preterm • Morbidity • Mortality • Monitoring • Management

KEY POINTS

- Late preterm infants are defined as those infants born between, and including, 34 weeks and 0 days and 36 weeks and 6 days.
- During the birth hospitalization, late preterm infants are at increased risk for morbidities such as respiratory distress, hypothermia, feeding difficulties, hyperbilirubinemia, and hypoglycemia.
- After discharge, late preterm infants are at increased risk for rehospitalization, mortality, and other morbidities, including neurologic, respiratory, developmental, and psychiatric/behavioral disorders.
- Monitoring for short-term and long-term morbidities is important in the late preterm population both during the birth hospitalization and into the future.

A significant increase in preterm births (those born <37 weeks of gestation) from 10.62% to 12.73% of all births occurred between 1990 and 2005.¹ Throughout this time, infants born later in the preterm period (variably referred to as near term, marginally preterm, mildly preterm, among other descriptors) continued to account for most, nearly three quarters, of these preterm births. To emphasize the importance of better optimizing the care of late preterm infants, along with supporting epidemiologic investigations into morbidity and mortality, the National Institute of Child Health and Human Development of the National Institutes of Health gathered a panel of experts to define this group. This panel defined infants born just before term as late preterm to emphasize their vulnerability and prematurity. Furthermore, late preterm was precisely defined as those born between, and including, the gestational ages of 34 weeks and 0 days (239 days) and 36 weeks and 6 days (259 days).²

Disclosure: The authors have no conflicts of interest to disclose.

Department of Neonatology, Indiana University School of Medicine, 699 Riley Hospital Drive, RR 208, Indianapolis, IN 46202, USA

* Corresponding author. Riley Research Room 208, 699 Riley Drive, Indianapolis, IN 46202.

E-mail address: wengle@iu.edu

Pediatr Clin N Am ■ (2018) ■–■

<https://doi.org/10.1016/j.pcl.2018.12.008>

0031-3955/18/© 2018 Elsevier Inc. All rights reserved.

pediatric.theclinics.com

This article describes epidemiologic trends in the rate of late preterm births and the associated risks for morbidities and mortality of being born late preterm. Furthermore, it discusses the importance of reducing the rate of late preterm births, monitoring and managing for short-term and long-term complications, and understanding factors that increase risk for morbidity and mortality in this subgroup of preterm infants.

Late preterm infants are a population with historically underappreciated increased risk for significant morbidity and mortality compared with term infants.^{2,3} In 2017, the rate of late preterm births in the United States increased for the third year in a row, up 5% since 2014, to 7.17% (Table 1).^{4,5} Late preterm births account for more than 70% of all preterm births, which equaled 279,382 births in 2016 alone (Fig. 1).⁴ Although morbidities occur in only a fraction of these infants, given the large number of late preterm births annually, these infants contribute to a large burden of care and cost to the health care and education systems compared with term infants.⁶

The recent increased rate of late preterm birth is concerning given the already large number that occur annually and prior successes in reducing such births between 2006 and 2014. The exact cause of this trend is unclear. Possible causes include the ongoing changes in maternal demographics (including increasing maternal age) and maternal health (including increasing rates of obesity and opioid use) that contribute to preterm birth.^{7–9} The increase in late preterm birth rate was observed across all maternal age groups, not just the older subgroups, so maternal age is less likely a new contributor.⁴ Past cited causes, including increased number of multiple gestations, high rate of elective inductions, or cesarean sections, have decreased in recent years and are less likely to contribute to this recent increase.⁴ Additional investigation is needed to clarify causation of this recent increase in late preterm birth rate and guide interventions to reverse this trend.

ACUTE MORBIDITIES DURING BIRTH HOSPITALIZATION

Late preterm infants are at risk for acute morbidities starting immediately after birth in the delivery room. Compared with term infants, late preterm infants more often require resuscitation (46% vs 28%; odds ratio [OR], 2.14; 95% confidence interval [CI], 1.88–2.44) including bag mask ventilation (14% vs 6%; OR, 2.61; 95% CI, 2.14–3.17).¹⁰ These increased rates of delivery room intervention persist even when controlling for variables such as cesarean delivery. The level of resuscitation required decreases

Year	Number	<28	28–31	32–33	34–36	37–38	39–40
2010	3,999,386	0.71	0.94	1.18	7.15	27.29	56.08
2011	3,953,590	0.70	0.93	1.18	6.99	26.09	57.51
2012	3,952,841	0.71	0.92	1.17	6.96	25.47	58.30
2013	3,932,181	0.70	0.92	1.17	6.83	24.81	58.85
2014	3,988,076	0.69	0.91	1.15	6.82	24.76	58.72
2015	3,978,487	0.68	0.91	1.17	6.87	24.99	58.47
2016	3,945,875	0.68	0.92	1.17	7.09	25.47	57.94
2017	3,853,472	—	—	—	7.17	—	—

Data from Martin JA, Hamilton BE, Osterman MJK, et al. Births: final data for 2016. US Department of Health and Human Services; 2018; and Hamilton BE, Martin JA, Osterman MJK, et al. Births: provisional data for 2017. Hyattsville (MD): National Center for Health Statistics; 2018.




Fig. 1. Breakdown of preterm births for 2016. (Data from Martin JA, Hamilton BE, Osterman MJK, et al. Births: final data for 2016. Hyattsville (MD): U. S. Department of Health and Human Services; 2018.)

with increasing gestational age between 34 and 39 weeks of gestation.¹¹ After delivery, late preterm infants are at increased risk of neonatal intensive care unit (NICU) admission.^{11,12} NICU admission criteria vary by institution but, even when comparing low-risk spontaneous late preterm delivery, late preterm infants continue to have an increased rate of NICU admission.¹² Reasons for NICU admission in this population include respiratory distress, hypothermia, feeding difficulties, hyperbilirubinemia, and hypoglycemia (Fig. 2).^{12,13}

Late preterm infants more often have symptoms of respiratory distress than term infants.¹³ With decreasing gestational age, there is an increase in respiratory distress syndrome, transient tachypnea of the newborn, pneumonia, apnea, and pneumothorax (Fig. 3).^{11,14} These respiratory morbidities lead to an increased need for respiratory support, including oxygen therapy, noninvasive ventilation, invasive ventilation, and surfactant replacement.^{11,14,15} Late preterm infants have structurally and functionally immature lungs with decreased surface area, decreased capacity for pulmonary fluid absorption at birth, and decreased surfactant




Fig. 2. Clinical morbidities noted during birth hospitalization from pooled data for late preterm and term patients. (Data from Refs. ^{11-13,77,78,90,91})




Fig. 3. Respiratory morbidities by gestational age. (Data from Teune MJ, Bakhuizen S, Gyamfi Bannerman C, et al. A systematic review of severe morbidity in infants born late preterm. *Am J Obstet Gynecol* 2011;205(4):374.e1-9.)

production.^{16–21} Antenatal betamethasone therapy has been shown to decrease the rates of respiratory morbidities in the late preterm population.²² Based on these observations, it is now recommended that women at risk of delivery between 34 0/7 weeks and 36 6/7 weeks of gestation receive a single course of corticosteroids.²³ Additional studies are needed to further understand the effect of this new recommendation on the rate of respiratory and other short-term and long-term morbidities in the late preterm population.²⁴

About 10% of late preterm infants experience temperature instability early during the birth hospitalization.¹³ Neonates in general rely on nonshivering thermogenesis to maintain body temperature.²⁵ Late preterm infants have lower amounts of brown adipose tissue and regulatory hormone concentrations compared with term infants, leading to a diminished ability to produce heat. Late preterm infants also have more heat loss than term infants because of lower amounts of white adipose tissue and higher surface area to mass ratio.^{3,19,25,26}

Feeding problems increase with decreasing gestational age; about 32% of late preterm infants compared with 7% of term infants have feeding difficulties during the birth hospitalization.¹³ Wang and colleagues¹³ found that poor feeding was the most common reason for discharge delay in late preterm infants. Feeding difficulties are caused by the physiologic immaturity of late preterm infants with decreased suck efficacy, abnormal sleep wake cycles, and problems with breathing and feeding coordination.^{27,28} These feeding difficulties lead to an increased need for supplemental feeds and an increased time to full oral feedings (median time to full oral feedings of 12 days at 34 weeks of gestation, 3 days at 35 weeks, and 2 days at 36 weeks).^{28,29} Feeding immaturity also complicates success with breastfeeding. However, because of the normal delay in full maternal milk supply, breastfeeding problems may initially be unrecognized until a larger amount of breastmilk is delivered by suckling.^{3,30,31} It is important to educate mothers, particularly first-time mothers, about such problems before discharging these infants. A higher rate of separation of late preterm infants from their mothers because of medical complications is also associated with a higher rate of difficulties with breastfeeding compared with term infants (eg, decreased maternal milk supply, increased time to oral feedings).^{29,32}

Feeding problems can increase the risk of physiologic morbidities such as hypoglycemia and hyperbilirubinemia in late preterm infants. Late preterm infants have lower glucose stores and glucose production compared with more mature infants because of hepatic enzymatic immaturity.^{3,33–35} Hepatic immaturity also contributes to more prolonged and clinically significant hyperbilirubinemia because of the lower concentrations of uridine diphosphate-glucuronosyltransferase, a key enzyme in the metabolism of hemoglobin.³⁶

As previously noted, late preterm infants are neurologically immature. At 34 weeks gestational age, a neonate's brain weighs 65% that of a term neonate.³⁷ Even at term postmenstrual gestational age, late preterm infants continue to have increased cerebral spinal fluid spaces, and decreased myelination, brain size, and biparietal diameter.³⁸ Such neurologic immaturity increases vulnerability to brain injury and long-term impairments in late preterm infants.³⁷

During the birth hospitalization, late preterm infants have a longer length of stay compared with more mature infants, with length of stay increasing with decreasing gestational age. During the initial birth hospitalization, infants born at 34 weeks' gestation stay on average 12.6 days compared with 6.1 days at 35 weeks birth gestational age and 3.8 days at 36 weeks birth gestational age (Fig. 4).³⁹ The most common factors contributing to increased length of stay in late preterm infants are feeding difficulties (75.9%), respiratory distress (30.8%), and jaundice (16.3%).¹³ In addition to gestational age, hospital practice variation regarding neonatal intensive care, intermediate nursery care, or well newborn nursery admission also affects the length of stay.⁴⁰ Such practice variation may lead to differences in outcomes after hospital discharge, given readmission rates vary in late preterm infants based on NICU admission and duration of birth hospitalization.⁴¹

Late preterm infants have an increased rate of mortality compared with term infants. In 2013, per 1000 live births, the infant mortality for gestational ages 34 to 36 weeks was 7.23 compared with 3.01 for 37 to 38 weeks and 1.85 for 39 to 40 weeks of gestation.⁴² Notably, mortality in late preterm infants varies with the reason for delivery.⁴³ Those infants born because of isolated spontaneous preterm labor have a neonatal mortality of 1.9 per 1000 live births. In comparison, late preterm infants born with no specific indication, a medical indication, or an obstetric indication have neonatal mortalities of 3.3, 3.8, and 8.6 per 1000 live births, respectively. Such different neonatal




Fig. 4. Mean length of stay by gestational age in weeks. (Data from Pulver LS, Denney JM, Silver RM, et al. Morbidity and discharge timing of late preterm newborns. *Clin Pediatr (Phila)* 2010;49(11):1061–7.)

mortalities emphasize the importance of causality of late preterm birth and associated intrauterine adversity for outcomes.

MORBIDITY IN LATE PRETERM INFANTS FOLLOWING THE BIRTH HOSPITALIZATION

Among infants of all gestational ages, readmission rates are highest among late preterm infants (<34 weeks, 3%; 34–36 weeks, 4.4%; >37 weeks, 2.0%).⁴⁴ A higher rate of readmission in late preterm infants correlates with a short duration of birth hospitalization (<4 days).⁴¹ Other risk factors for readmission include any labor or delivery complication, primigravid mother, breastfeeding, and Asian/Pacific Islander ethnicity.⁴⁵ Common reasons for readmission include jaundice, feeding difficulties, respiratory distress/apnea, and infection.^{41,44,46,47} Jaundice is also a frequent presenting complaint for emergency department visits among neonates, with late preterm infants, especially those born at 36 weeks of gestation, presenting more often than infants of other gestational ages.⁴⁸

Later in life, long-term morbidities encountered by late preterm infants include neurodevelopmental, respiratory, and medical illnesses. Mortality during young adulthood is also higher in late preterm infants than in term infants, with rates of 0.65 per 1000 person years versus 0.46 per 1000 person years, respectively (hazard ratio, 1.31; 95% CI, 1.13–1.50).⁴⁹ Late preterm infants have increased rates of developmental delay and cerebral palsy compared with term infants.^{50–52} For example, rates of developmental delay/cognitive dysfunction and cerebral palsy at 34 weeks of gestation are 12 per 1000 children and 9 per 1000 children, respectively; at 38 to 41 weeks of gestation, comparable rates are 9 per 1000 and 3 per 1000 children, respectively.^{50,52} Partly because of higher rates of neurodevelopmental impairments, more late preterm infants are enrolled in early intervention programs compared with term infants (23.5% vs 11.9%, respectively).⁵³

Late preterm infants are also more likely to have behavioral problems, school difficulties, and autism.^{51,54–56} Behavioral problems are more common in those diagnosed with developmental delays.⁵⁴ However, an increased risk of attention-deficit/hyperactivity disorder has not been consistently found.^{57,58} A higher rate of psychiatric hospitalization among adolescents born late preterm has been reported.^{59,60} Neurodevelopmental disabilities and impairments are also found in adults born late preterm.^{60,61} In a Norwegian cohort of adults aged 20 to 36 years, former late preterm infants had a higher rate of cognitive dysfunction, schizophrenia, disorders of psychological development, behavior and emotional disorders, and disabilities affecting their capacity to perform work.⁶⁰ Following late preterm birth, low neurocognitive test scores have been reported in adults in their late 60s.⁶² Furthermore, late preterm infants have lower socioeconomic attainment in late adulthood with lower levels of education and income.^{61,63} Other medical diagnoses, including diabetes treated with oral medications or insulin and hypertension, are also more prevalent in former late preterm infants.^{64,65}

After the birth hospitalization, late preterm infants continue to be at increased risk of respiratory morbidities. In the first 2 years after birth, previously late preterm infants experience increased morbidity with common childhood illnesses. Late preterm infants have a higher likelihood of needing hospitalization with respiratory syncytial virus infection compared with term infants (2.5% vs 1.3%, respectively).⁶⁶ Those presenting with respiratory illnesses also have an increased need for intensive care and longer durations of hospital stay than more mature infants.⁶⁷ At 18 months of age there is an increased rate of persistent asthma (adjusted OR [aOR], 1.68; 95% CI, 1.01–2.80), need for inhaled corticosteroids (aOR, 1.66; 95% CI, 1.20–2.29), and number of urgent

visits for respiratory disorders (incident rate ratio, 1.45; 95% CI, 1.25–1.68).⁶⁸ During late childhood and adolescence, poor pulmonary function tests are found in late preterm compared with term infants.^{69,70} To control for environmental confounders, especially smoke exposure, contributing to respiratory morbidity, Todisco and colleagues⁷⁰ compared late preterm infants with their term siblings and found increased residual volume and residual volume to total lung capacity ratio. Therefore, even when accounting for postnatal exposures, respiratory outcome differences persist, suggesting that intrinsic pulmonary abnormalities are present during childhood and beyond in former late preterm infants.

RISK FACTORS FOR MORBIDITY AND MORTALITY IN LATE PRETERM INFANTS

Given the increased rate of morbidity and mortality of late preterm infants, close monitoring for social, neurodevelopmental, and medical disabilities beyond the neonatal period is warranted. However, most late preterm infants survive, grow, and develop without identifiable medical or social difficulties. This lack of morbidities and mortality may reflect an absence of complications or evaluation at an age when problems have not yet presented.^{71–76} Thus, an important question is: what factors are associated with development of complications of late prematurity? The answer is likely multifactorial and may include combinations of influences that include physical and physiologic immaturity, genetic endowment, fetal environmental exposures and stressors, congenital anomalies, infections, medical disorders, and complications associated with any of the preceding factors, along with the postnatal environment.^{71,77,78}

In the United States, congenital anomalies are the leading cause of infant mortality (129 per 1000 live births) and second leading cause of neonatal mortality after prematurity-associated complications (92.1 per 1000 live births).⁷⁹ In a Swedish cohort of 14,030 late preterm infants, 41% of late preterm neonatal deaths (1.14 per 1000 live births) were attributed to congenital anomalies.⁷⁷ Comparing the most commonly noted indications for late preterm delivery, major congenital anomaly is associated with the highest rate of neonatal (107.9 per 1000 live births) and infant (140.7 per 1000 live births) mortality.⁴³

The cause of late preterm birth may be associated with an adverse intrauterine environment and is an important determinant of fetal outcomes, including neonatal and infant mortality; acute birth hospitalization conditions; and long-term neurodevelopmental, medical, and social health.^{43,77} In addition to congenital anomalies, other causes of late preterm delivery, categorized as obstetric, medical, no recorded indication, and isolated spontaneous labor, are associated with infant mortalities of 13.3, 7.0, 6.8, and 4.8 per 1000 live births, respectively.⁴³ Maternal factors affecting late preterm delivery and neonatal and infant outcomes include maternal race, marital status, age, smoking status, health (including diabetes and hypertensive disorders), diet, income, and educational levels.^{42,71,76–78,80,81} Pregnancy, labor, and delivery complications, including multiple gestation, intrauterine growth restriction, antepartum hemorrhage, abruption, preterm prelabor rupture of membranes, and chorioamnionitis, also occur at higher rates in late preterm pregnancies.^{77,78,81,82} Such complications contribute to premature birth and likely morbidity and mortality of late preterm infants.

Physiologic and structural immaturity are important fetal and neonatal contributions to the risks of morbidity and mortality in late preterm infants.^{3,30,51,74} The rates of neonatal intensive care admission and acute birth hospital morbidities increase with decreasing gestational age, consistent with a major impact of immaturity on such outcomes.^{11,13,77} Immaturity also adversely affects neurodevelopmental outcomes with

decreasing gestational age in neonates without other genetic, metabolic, or neurologic disorders.^{51,72} Across the gestational age spectrum from 32 to 41 weeks, the risk for neurodevelopmental delay compared with a term cohort is much higher in the younger gestational age groups. This risk of developmental delay for the moderately preterm (32 0/7 to 33 6/7 weeks), late preterm (34 0/7 to 36 6/7 weeks), and early term (37 0/7 to 38 6/7 weeks) infants is 3-fold (aOR, 3.01; 95% CI, 1.59–5.71), 2.6-fold (aOR, 2.58; 95% CI, 1.66–4.01), and 1.6-fold (aOR, 1.56; 95% CI, 1.19–2.06), respectively, compared with those born at 39 0/7 to 41 6/7 weeks gestational age.⁷² In a cohort of 22,552 healthy late preterm infants (ie, discharged from birth hospitalization within 3 days of birth), 4.24% of late preterm and 2.96% of full-term infants were diagnosed with developmental delay or disability (adjusted RR, 1.36; 95% CI, 1.29–1.43).⁵¹ In this same cohort, referral for prekindergarten services, exceptional student education, retention in kindergarten, and suspension in kindergarten were all significantly increased in the late preterm group.

In contrast with the association of immaturity with poor neurodevelopmental outcomes in nonanomalous late preterm versus term infants, conflicting evidence has been reported for an association with specific outcomes during early childhood, such as intellectual disability (intelligence quotient <70), overall developmental score, problem solving, attention and behavior disorders, and learning disabilities.^{57,71,73–75,83} Such contrasting results may be associated with absence of true difference in outcomes, variations in sensitivity of study methods to differentiate outcomes at some ages, methodologic issues, and population sample size limitations. Despite some contradictory evidence, the preponderance of information, including differences observed through adulthood, support an important contribution of immaturity on neurodevelopmental, educational, social, medical, and behavioral outcomes of infants born late preterm.^{3,30,51,60–63,74}

In addition to immaturity, other fetal/neonatal factors, including sex and birth weight, specifically infants small for gestational age, are associated with late preterm delivery in addition to other outcomes, including mortality.^{42,77,82} Late preterm infants who are also small for gestational age have been shown to have an increased risk for poor neurocognitive intelligence test scores in young adulthood (full-scale intelligence quotient score, -11.84 ; 95% CI, -18.33 to -5.36 compared with term appropriate for gestational age infants) and cerebral palsy in childhood (pooled OR, 3.47; 95% CI, 1.29–9.31), indicating the lasting impact of intrauterine growth restriction on this population.^{71,84}

The morbidity and mortality risks found in late preterm infants compared with term infants are numerous. Adverse outcomes throughout the lifespan of infants born late preterm are the result of an interplay between immaturity and a host of unique pathological and social determinants that also contribute to poor outcomes in infants born at other gestational ages.

COSTS ATTRIBUTABLE TO LATE PRETERM BIRTHS

The costs, including emotional and financial, for families and society attributable to consequences of late preterm birth are extensive.^{85–87} Although the percentage of late preterm infants without congenital anomalies that have long-term impairments is low compared with extreme and moderate preterm infants, the large absolute number of late preterm births (in 2016, 279, 382 late preterm vs 108,836 extreme and moderate preterm infants) contributes to the large number of children in need of special care and support services.⁴ For example, using population-based data from Finland, where 0.55% of late preterm and 0.85% of combined extremely and moderately

Table 2 Sample management and monitoring guideline for late preterm infants during the birth hospitalization	
Delivery Room ^{30,88,89,92}	<ul style="list-style-type: none"> • Monitor for signs of respiratory distress • NRP Guidelines followed if resuscitation is needed • Thermoregulation measures: hat and skin to skin or radiant warmer • Vital signs (heart rate, respiratory rate, and temperature) measured within the first hour of birth
NICU Admission Decision ³⁰	<ul style="list-style-type: none"> • Significant resuscitation in the delivery room (ie, positive pressure ventilation, chest compressions, medications) or clinical concerns, observe patient with continuous vital sign monitoring for at least 6 h • Institution-specific guideline for gestational age and weight parameters for NICU admission considered when determining admission placement
Thermoregulation ^{30,88,89}	<ul style="list-style-type: none"> • Temperature measurement every hour for the first 6 h and then minimum of every 6 h until discharge • If temperature is < 36.0°C, swaddle infant and cover head with a hat. In 30 min, if temperature remains <36.0°C, place infant under radiant warmer. If still <36.0°C despite radiant warmer, admit infant to higher level of care, if in Well Baby Nursery, for further evaluation and management.
Respiratory Distress ^{30,89}	<ul style="list-style-type: none"> • During the first 2 h after delivery infants may have abnormal respiratory rates (rate 25–100 breaths per minute). Normalization is expected after this transitional period (respiratory rate 40–60 breaths per minute). • Monitor for signs of respiratory distress throughout the birth hospitalization. • If signs of respiratory distress arise, institute cardiorespiratory monitoring with consideration to move infant to higher level of care. • Apnea with pause in breathing >20 s also requires closer cardiorespiratory monitoring and transfer to higher level of care.
Feeding ^{3,30,88,89}	<ul style="list-style-type: none"> • All aspects of feedings monitored including latch, frequency, volume, and duration. • Daily weights monitored. • Optimize feeding by having a health care professional observe at least one feeding each shift. If the infant is breastfeeding, this may be performed by a lactation consultant. • If feeding difficulty noted, early consultation with speech therapy or occupational therapy. • If excessive weight loss (more than 3% per day of age or more than 7% of birth weight total) consider further evaluation for dehydration and possible supplementation. • If the infant is unable to maintain adequate oral intake other forms of feeding, such as gavage feedings are recommended.
Hypoglycemia ⁹³	<ul style="list-style-type: none"> • <i>Birth to 4 h of age:</i> Encourage feeding within 1 h of birth and measure glucose 30 min after feeding. If glucose is < 25 mg/dL refeeding is recommended. If glucose still <25 mg/dL 1 hour after feeding IV glucose is recommended. If glucose is 25–40 mg/dL refeeding or IV glucose is recommended. • <i>From 4 h to 24 h of age:</i> Glucose measurement is recommended before each feeding. If < 35 mg/dL feed, and recheck in 1 h. If still <35 mg/dL IV glucose is recommended. If glucose 35–45 mg/dL encourage feeding or consider IV glucose. • If symptomatic hypoglycemia, give IV glucose.

(continued on next page)

Table 2
(continued)

Hyperbilirubinemia ^{94,95}	<ul style="list-style-type: none"> ● Assess risk factors for hyperbilirubinemia, including ABO incompatibility and presence of maternal antibody for each infant. ● If maternal blood type is Rh negative or O Rh positive, determine infant blood type. ● Monitor clinical signs of jaundice regularly. ● Within the first 24 h, obtain transcutaneous or serum total bilirubin. ● For infants >35 wk of age, treat hyperbilirubinemia with phototherapy per the AAP guidelines. ● For infants <35 wk, initiate phototherapy at a lower threshold. Because of inadequate evidence regarding the treatment threshold in this group, each institution should establish their own guidelines.
-------------------------------------	---

Abbreviation: NRP, neonatal resuscitation program.

preterm infants have intellectual disability, late preterm births would contribute 1537 children with intellectual disability compared with 926 extremely and moderately preterm births in 2016 alone.⁸³

MONITORING AND MANAGEMENT OF LATE PRETERM INFANTS

Given the significant morbidities encountered by late preterm infants, this population warrants increased observation and close monitoring during birth hospitalization, soon after discharge, and throughout life. Understanding the morbidities and the time frame during which they may occur can guide health care professionals to care for this at-risk population. Monitoring for these morbidities and ensuring adequate care of late preterm infants will help to decrease rates of negative outcomes.

During the birth hospitalization, as previously mentioned, late preterm infants are at risk for respiratory distress, hypothermia, hypoglycemia, hyperbilirubinemia, and feeding difficulties.^{12,13} **Table 2** provides a sample guideline for monitoring and managing these morbidities. Clinicians who care for late preterm infants during their birth hospitalization should consider setting guidelines to care for these infants, because

Table 3
Discharge criteria recommendations for late preterm infants

<ul style="list-style-type: none"> ● In general, a minimum of 48 h of age unless necessary competencies established beforehand ● 12–24 h of stable vital signs (heart rate, respiratory rate, temperature) ● Physical examination without abnormalities for which ongoing hospital management is necessary ● Adequate feeding for at least 24 h with at least 2 feedings observed by health care professionals ● If weight loss more than 3% per day, or 7% total, at time of discharge, consider dehydration evaluation and/or supplementation before release 	<ul style="list-style-type: none"> ● Passage of at least 1 stool ● Bilirubin level (transcutaneous or serum) documented and does not require intervention ● All state-mandated screening evaluations performed (newborn metabolic, hearing, congenital heart screens) ● Car seat study passed if applicable ● Family and social situation assessed and appropriate for discharge ● Outpatient follow-up scheduled within 24–48 h of discharge
---	---

Data from Refs. ^{3,30,88,89}

variation in practice effects the length of stay, care received, and potentially outcomes after discharge (eg, readmission rates).⁴⁰ Before discharge, it is recommended that specific goals be met (**Table 3**). Other considerations before discharge include the parents' ability to provide care for the infant at the bedside. This ability of parents to care for their infants during the birth hospitalization has been linked to outcomes and future maternal infant interactions after discharge.⁸⁷

After birth hospitalization, it is recommended that late preterm infants be followed closely by their primary care physicians. Initial follow-up is recommended within 24 to 48 hours to evaluate the infant for feeding issues, jaundice, and other concerns.^{30,88,89} Frequency of follow-up can be increased if issues arise, until the infant is clearly thriving and developing.⁸⁹ It is important for previously late preterm infants to have an established primary care physician.⁸⁹ These infants should receive all recommended screens throughout childhood, with care taken to diagnose disabilities or medical disorders as early as possible. If a problem is identified, such as a developmental delay, referral for the appropriate intervention or therapies should be made as early as possible. Parents should be informed of the increased risk of long-term morbidities associated with prematurity, to emphasize the importance of continued timely follow-up of late preterm infants.

SUMMARY

Late preterm infants are at increased risk of significant morbidity and mortality. During birth hospitalization, these morbidities include respiratory distress, hypothermia, feeding difficulties, hypoglycemia, and hyperbilirubinemia.^{12,13} After discharge, these infants are at increased risk of rehospitalization because of feeding difficulties and jaundice, especially with shorter birth hospitalization.^{41,43,44,46,47} Risks for morbidity and mortality continue throughout life for late preterm infants. This persistence emphasizes the need for well-defined long-term follow-up.

REFERENCES

1. Martin JA, Hamilton BE, Ventura SJ, et al. Births: final data for 2010. *Natl Vital Stat Rep* 2012;61(1):1–72.
2. Raju TN, Higgins RD, Stark AR, et al. Optimizing care and outcome for late-preterm (near-term) infants: a summary of the workshop sponsored by the National Institute of Child Health and Human Development. *Pediatrics* 2006; 118(3):1207–14.
3. Engle WA, Tomashek KM, Wallman C. “Late-preterm” infants: a population at risk. *Pediatrics* 2007;120(6):1390–401.
4. Martin JA, Hamilton BE, Osterman MJK, et al. Births: final data for 2016. Hyattsville (MD): US Department of Health and Human Services; 2018.
5. Hamilton BE, Martin JA, Osterman MJK, et al. Births: provisional data for 2017. Hyattsville (MD): National Center for Health Statistics; 2018.
6. McLaurin KK, Hall CB, Jackson EA, et al. Persistence of morbidity and cost differences between late-preterm and term infants during the first year of life. *Pediatrics* 2009;123(2):653–9.
7. Deputy NP, Dub B, Sharma AJ. Prevalence and trends in prepregnancy normal weight - 48 states, New York City, and District of Columbia, 2011-2015. *MMWR Morb Mortal Wkly Rep* 2018;66(5152):1402–7.
8. Martin JA, Hamilton BE, Osterman MJK. Births in the United States, 2016. Hyattsville (MD): National Center for Health Statistics; 2017.

9. Norgaard M, Nielsson MS, Heide-Jorgensen U. Birth and neonatal outcomes following opioid use in pregnancy: a Danish population-based study. *Subst Abuse* 2015;9(Suppl 2):5–11.
10. de Almeida MF, Guinsburg R, da Costa JO, et al. Resuscitative procedures at birth in late preterm infants. *J Perinatol* 2007;27(12):761–5.
11. Hibbard JU, Wilkins I, Sun L, et al. Respiratory morbidity in late preterm births. *JAMA* 2010;304(4):419–25.
12. Melamed N, Klinger G, Tenenbaum-Gavish K, et al. Short-term neonatal outcome in low-risk, spontaneous, singleton, late preterm deliveries. *Obstet Gynecol* 2009; 114(2 Pt 1):253–60.
13. Wang ML, Dorer DJ, Fleming MP, et al. Clinical outcomes of near-term infants. *Pediatrics* 2004;114(2):372–6.
14. Teune MJ, Bakhuizen S, Gyamfi Bannerman C, et al. A systematic review of severe morbidity in infants born late preterm. *Am J Obstet Gynecol* 2011;205(4): 374.e1-9.
15. Gouyon JB, Vintejoux A, Sagot P, et al. Neonatal outcome associated with singleton birth at 34-41 weeks of gestation. *Int J Epidemiol* 2010;39(3):769–76.
16. Bland RD. Loss of liquid from the lung lumen in labor: more than a simple “squeeze”. *Am J Physiol Lung Cell Mol Physiol* 2001;280(4):L602–5.
17. Kallapur SG, Jobe AH. Lung development and maturation. In: Martin RJ, Fanaroff AA, Walsh MC, editors. *Fanaroff and Martin's neonatal perinatal medicine*. Philadelphia: Elsevier Saunders; 2015. p. 1042–59.
18. Pike KC, Lucas JS. Respiratory consequences of late preterm birth. *Paediatr Respir Rev* 2015;16(3):182–8.
19. Ramachandrapa A, Jain L. The late preterm infant. In: Martin R, Fanaroff A, Walsh M, editors. *Fanaroff and Martin's neonatal-perinatal medicine*. Philadelphia: Elsevier Saunders; 2015. p. 577–91.
20. Rozycki HJ, Hendricks-Munoz KD. Structure and development of alveolar epithelial cells. In: Polin RA, Abman SH, Rowitch DH, et al, editors. *Fetal and neonatal physiology*, vol. 1, 5th edition. Philadelphia: Elsevier; 2017. p. 809–13.
21. Smith DE, Otulakowski G, Yeger H, et al. Epithelial Na(+) channel (ENaC) expression in the developing normal and abnormal human perinatal lung. *Am J Respir Crit Care Med* 2000;161(4 Pt 1):1322–31.
22. Gyamfi-Bannerman C, Thom EA, Blackwell SC, et al. Antenatal betamethasone for women at risk for late preterm delivery. *N Engl J Med* 2016;374(14):1311–20.
23. Committee on Obstetric Practice. Committee opinion no. 713: antenatal corticosteroid therapy for fetal maturation. *Obstet Gynecol* 2017;130(2):e102–9.
24. Kaempf JW, Suresh G. Antenatal corticosteroids for the late preterm infant and agnology. *J Perinatol* 2017;37(12):1265–7.
25. Sahni R. Temperature control in newborn infants. In: Polin RA, Abman SH, Rowitch DH, et al, editors. *Fetal and neonatal physiology*, vol. 1, 5th edition. Philadelphia: Elsevier; 2017. p. 459–82.
26. Symonds ME, Mostyn A, Pearce S, et al. Endocrine and nutritional regulation of fetal adipose tissue development. *J Endocrinol* 2003;179(3):293–9.
27. Margolis KG, Picoraro JA. Development of gastrointestinal motility. In: Polin RA, Abman SH, Rowitch DH, et al, editors. *Fetal and neonatal physiology*, vol. 1, 5th edition. Philadelphia: Elsevier; 2017. p. 881–8.
28. Mizuno K, Ueda A. The maturation and coordination of sucking, swallowing, and respiration in preterm infants. *J Pediatr* 2003;142(1):36–40.
29. Jackson BN, Kelly BN, McCann CM, et al. Predictors of the time to attain full oral feeding in late preterm infants. *Acta Paediatr* 2016;105(1):e1–6.

30. Engle WA. Morbidity and mortality in late preterm and early term newborns: a continuum. *Clin Perinatol* 2011;38(3):493–516.
31. McManaman JL. Physiology of lactation. In: Polin RA, Abman SH, Rowitch DH, et al, editors. *Fetal and neonatal physiology*, vol. 1, 5th edition. Philadelphia: Elsevier; 2017. p. 281–7.
32. Briere CE, Lucas R, McGrath JM, et al. Establishing breastfeeding with the late preterm infant in the NICU. *J Obstet Gynecol Neonatal Nurs* 2015;44(1):102–13 [quiz: E101–2].
33. Garg M, Devaskar SU. Glucose metabolism in the late preterm infant. *Clin Perinatol* 2006;33(4):853–70.
34. Hume R, Burchell A. Abnormal expression of glucose-6-phosphatase in preterm infants. *Arch Dis Child* 1993;68(2):202–4.
35. Stanley CA, Rozance PJ, Thornton PS, et al. Re-evaluating “transitional neonatal hypoglycemia”: mechanism and implications for management. *J Pediatr* 2015; 166(6):1520–5.e1.
36. Kawade N, Onishi S. The prenatal and postnatal development of UDP-glucuronyltransferase activity towards bilirubin and the effect of premature birth on this activity in the human liver. *Biochem J* 1981;196(1):257–60.
37. Kinney HC. The near-term (late preterm) human brain and risk for periventricular leukomalacia: a review. *Semin Perinatol* 2006;30(2):81–8.
38. Walsh JM, Doyle LW, Anderson PJ, et al. Moderate and late preterm birth: effect on brain size and maturation at term-equivalent age. *Radiology* 2014;273(1): 232–40.
39. Pulver LS, Denney JM, Silver RM, et al. Morbidity and discharge timing of late preterm newborns. *Clin Pediatr* 2010;49(11):1061–7.
40. Aliaga S, Boggess K, Ivester TS, et al. Influence of neonatal practice variation on outcomes of late preterm birth. *Am J Perinatol* 2014;31(8):659–66.
41. Escobar GJ, Joffe S, Gardner MN, et al. Rehospitalization in the first two weeks after discharge from the neonatal intensive care unit. *Pediatrics* 1999;104(1):e2.
42. Mathews TJ, MacDorman MF, Thoma ME. Infant mortality statistics from the 2012 period linked birth/infant death data set. *Natl Vital Stat Rep* 2015;64(9):1–30.
43. Reddy UM, Ko CW, Raju TN, et al. Delivery indications at late-preterm gestations and infant mortality rates in the United States. *Pediatrics* 2009;124(1):234–40.
44. Escobar GJ, Greene JD, Hulac P, et al. Rehospitalisation after birth hospitalisation: patterns among infants of all gestations. *Arch Dis Child* 2005;90(2):125–31.
45. Shapiro-Mendoza CK, Tomashek KM, Kotelchuck M, et al. Risk factors for neonatal morbidity and mortality among “healthy,” late preterm newborns. *Semin Perinatol* 2006;30(2):54–60.
46. Tomashek KM, Shapiro-Mendoza CK, Weiss J, et al. Early discharge among late preterm and term newborns and risk of neonatal morbidity. *Semin Perinatol* 2006; 30(2):61–8.
47. Young PC, Korgenski K, Buchi KF. Early readmission of newborns in a large health care system. *Pediatrics* 2013;131(5):e1538–44.
48. Jain S, Cheng J. Emergency department visits and rehospitalizations in late preterm infants. *Clin Perinatol* 2006;33(4):935–45 [abstract: xi].
49. Crump C, Winkleby MA, Sundquist J, et al. Risk of asthma in young adults who were born preterm: a Swedish national cohort study. *Pediatrics* 2011;127(4): e913–20.
50. Hirvonen M, Ojala R, Korhonen P, et al. Cerebral palsy among children born moderately and late preterm. *Pediatrics* 2014;134(6):e1584–93.

51. Morse SB, Zheng H, Tang Y, et al. Early school-age outcomes of late preterm infants. *Pediatrics* 2009;123(4):e622–9.
52. Petrini JR, Dias T, McCormick MC, et al. Increased risk of adverse neurological development for late preterm infants. *J Pediatr* 2009;154(2):169–76.
53. Shapiro-Mendoza C, Kotelchuck M, Barfield W, et al. Enrollment in early intervention programs among infants born late preterm, early term, and term. *Pediatrics* 2013;132(1):e61–9.
54. Potijk MR, de Winter AF, Bos AF, et al. Co-occurrence of developmental and behavioural problems in moderate to late preterm-born children. *Arch Dis Child* 2016;101(3):217–22.
55. Talge NM, Holzman C, Wang J, et al. Late-preterm birth and its association with cognitive and socioemotional outcomes at 6 years of age. *Pediatrics* 2010;126(6):1124–31.
56. Chan E, Quigley MA. School performance at age 7 years in late preterm and early term birth: a cohort study. *Arch Dis Child Fetal Neonatal Ed* 2014;99(6):F451–7.
57. Harris MN, Voigt RG, Barbaresi WJ, et al. ADHD and learning disabilities in former late preterm infants: a population-based birth cohort. *Pediatrics* 2013;132(3):e630–6.
58. Linnet KM, Wisborg K, Agerbo E, et al. Gestational age, birth weight, and the risk of hyperkinetic disorder. *Arch Dis Child* 2006;91(8):655–60.
59. Lindstrom K, Lindblad F, Hjern A. Psychiatric morbidity in adolescents and young adults born preterm: a Swedish national cohort study. *Pediatrics* 2009;123(1):e47–53.
60. Moster D, Lie RT, Markestad T. Long-term medical and social consequences of preterm birth. *N Engl J Med* 2008;359(3):262–73.
61. Lindstrom K, Winbladh B, Haglund B, et al. Preterm infants as young adults: a Swedish national cohort study. *Pediatrics* 2007;120(1):70–7.
62. Heinonen K, Eriksson JG, Lahti J, et al. Late preterm birth and neurocognitive performance in late adulthood: a birth cohort study. *Pediatrics* 2015;135(4):e818–25.
63. Heinonen K, Eriksson JG, Kajantie E, et al. Late-preterm birth and lifetime socio-economic attainments: the Helsinki Birth Cohort Study. *Pediatrics* 2013;132(4):647–55.
64. Crump C, Winkleby MA, Sundquist K, et al. Risk of diabetes among young adults born preterm in Sweden. *Diabetes Care* 2011;34(5):1109–13.
65. Gunay F, Alpay H, Gokce I, et al. Is late-preterm birth a risk factor for hypertension in childhood? *Eur J Pediatr* 2014;173(6):751–6.
66. Helfrich AM, Nylund CM, Eberly MD, et al. Healthy late-preterm infants born 33–36+6 weeks gestational age have higher risk for respiratory syncytial virus hospitalization. *Early Hum Dev* 2015;91(9):541–6.
67. Gunville CF, Sontag MK, Stratton KA, et al. Scope and impact of early and late preterm infants admitted to the PICU with respiratory illness. *J Pediatr* 2010;157(2):209–14.e1.
68. Goyal NK, Fiks AG, Lorch SA. Association of late-preterm birth with asthma in young children: practice-based study. *Pediatrics* 2011;128(4):e830–8.
69. Kotecha SJ, Watkins WJ, Paranjothy S, et al. Effect of late preterm birth on longitudinal lung spirometry in school age children and adolescents. *Thorax* 2012;67(1):54–61.
70. Todisco T, de Benedictis FM, Iannacci L, et al. Mild prematurity and respiratory functions. *Eur J Pediatr* 1993;152(1):55–8.

71. Heinonen K, Lahti J, Sammallahti S, et al. Neurocognitive outcome in young adults born late-preterm. *Dev Med Child Neurol* 2018;60(3):267–74.
72. Schonhaut L, Armijo I, Perez M. Gestational age and developmental risk in moderately and late preterm and early term infants. *Pediatrics* 2015;135(4):e835–41.
73. Gurka MJ, LoCasale-Crouch J, Blackman JA. Long-term cognition, achievement, socioemotional, and behavioral development of healthy late-preterm infants. *Arch Pediatr Adolesc Med* 2010;164(6):525–32.
74. Hornman J, de Winter AF, Kerstjens JM, et al. Stability of developmental problems after school entry of moderately-late preterm and early preterm-born children. *J Pediatr* 2017;187:73–9.
75. Shah P, Kaciroti N, Richards B, et al. Developmental outcomes of late preterm infants from infancy to kindergarten. *Pediatrics* 2016;138(2) [pii:e20153496].
76. Woythaler M, McCormick MC, Mao WY, et al. Late preterm infants and neurodevelopmental outcomes at kindergarten. *Pediatrics* 2015;136(3):424–31.
77. Bonnevier A, Brodzski J, Bjorklund LJ, et al. Underlying maternal and pregnancy-related conditions account for a substantial proportion of neonatal morbidity in late preterm infants. *Acta Paediatr* 2018;107:1521–8.
78. Dimitriou G, Fouzas S, Georgakis V, et al. Determinants of morbidity in late preterm infants. *Early Hum Dev* 2010;86(9):587–91.
79. Ely DM, Hoyert DL. Differences between rural and urban areas in mortality rates for the leading causes of infant death: United States, 2013-2015. *NCHS Data Brief* 2018;(300):1–8.
80. Smith LK, Draper ES, Evans TA, et al. Associations between late and moderately preterm birth and smoking, alcohol, drug use and diet: a population-based case-cohort study. *Arch Dis Child Fetal Neonatal Ed* 2015;100(6):F486–91.
81. Manuck TA, Rice MM, Bailit JL, et al. Preterm neonatal morbidity and mortality by gestational age: a contemporary cohort. *Am J Obstet Gynecol* 2016;215(1):103.e1–14.
82. Mathiasen R, Hansen BM, Andersen AM, et al. Gestational age and basic school achievements: a national follow-up study in Denmark. *Pediatrics* 2010;126(6):e1553–61.
83. Hirvonen M, Ojala R, Korhonen P, et al. Intellectual disability in children aged less than seven years born moderately and late preterm compared with very preterm and term-born children - a nationwide birth cohort study. *J Intellect Disabil Res* 2017;61(11):1034–54.
84. Zhao M, Dai H, Deng Y, et al. SGA as a risk factor for cerebral palsy in moderate to late preterm infants: a system review and meta-analysis. *Sci Rep* 2016;6:38853.
85. Haataja P, Korhonen P, Ojala R, et al. Hospital admissions for lower respiratory tract infections in children born moderately/late preterm. *Pediatr Pulmonol* 2018;53(2):209–17.
86. Isayama T, Lewis-Mikhael AM, O'Reilly D, et al. Health services use by late preterm and term infants from infancy to adulthood: a meta-analysis. *Pediatrics* 2017;140(1) [pii:e20170266].
87. Premji SS, Pana G, Currie G, et al. Mother's level of confidence in caring for her late preterm infant: a mixed methods study. *J Clin Nurs* 2018;27(5–6):e1120–33.
88. Kilpatrick S, Papile L, Macones G. Guidelines for perinatal care. 8th edition. Elk Grove Village, IL and Washington, DC: American Academy of Pediatrics and American College of Obstetricians and Gynecologists; 2017.

89. Phillips RM, Goldstein M, Houglan K, et al. Multidisciplinary guidelines for the care of late preterm infants. *J Perinatol* 2013;33(Suppl 2):S5–22.
90. Boyle EM, Johnson S, Manktelow B, et al. Neonatal outcomes and delivery of care for infants born late preterm or moderately preterm: a prospective population-based study. *Arch Dis Child Fetal Neonatal Ed* 2015;100(6):F479–85.
91. Sarici SU, Serdar MA, Korkmaz A, et al. Incidence, course, and prediction of hyperbilirubinemia in near-term and term newborns. *Pediatrics* 2004;113(4):775–80.
92. Weiner G, Zaickin J. Textbook of neonatal resuscitation. 7th edition. Elk Grove Village (IL): American Academy of Pediatrics; 2016.
93. Adamkin DH. Postnatal glucose homeostasis in late-preterm and term infants. *Pediatrics* 2011;127(3):575–9.
94. American Academy of Pediatrics Subcommittee on Hyperbilirubinemia. Management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation. *Pediatrics* 2004;114(1):297–316.
95. Maisels MJ, Watchko JF, Bhutani VK, et al. An approach to the management of hyperbilirubinemia in the preterm infant less than 35 weeks of gestation. *J Perinatol* 2012;32(9):660–4.